

ON THE DIFFERENCE EQUATION OF THE POINCARÉ TYPE

L.A. Gutnik

Dedicated to the memory of Professor A.O. Gelfond.

Abstract. In the paper are proved theorems, which amplify the results of my paper "On the difference equation of Poincaré type (Part 3)", Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2004, 09, 1 – 34.

Table of contents

- §0. Foreword.
- §1. Some preparatory results.
- §2. Proof of the theorem 8.
- §3. Proof of the Theorem 9.
- §4. Proof of the Theorem 10.
- §5. The case of the general difference equation of the Poincaré type.

§0. Foreword.

In [77]: was proved the following

Theorem 7. *Let $s \in \mathbb{N} - 1, n \in \mathbb{N}$,*

$$a_i^{\sim} \in \mathbb{C}, a_i(\nu) \in \mathbb{C},$$

$$(1) \quad a_n(\nu) = 1, a_i(\nu) - a_i^{\sim} = O(1/(\nu + 1))$$

for $\nu \in \mathbb{N} - 1$ and $i = 0, \dots, n$. Let us consider the following difference equation

$$(2) \quad \sum_{k=0}^n a_k(\nu) y(\nu + k) = 0,$$

where $\nu \in \mathbb{N} - 1$.

For $m \in \mathbb{N}$ let V_m denotes the linear over $\mathbb{C}$ space of solutions $y = y(\nu)$ of the equation

$$(3) \quad \sum_{k=0}^n a_k(\nu) y(\nu + k) = 0,$$

where $\nu \in m + \mathbb{N} - 1$. Let the absolute values of all the roots of the characteristic polynomial

$$(4) \quad T(z) = \sum_{k=0}^n a_k^{\sim} z^k$$

are among the numbers $\{\rho_i: 1 \leq i \leq 1 + s\}$ such that $\rho_{s+1} = 0$ and $\rho_j < \rho_i$ for $1 \leq i < j \leq s + 1$. Let e_i and k_i denote respectively the sum and the maximum of the multiplicities of those roots, whose absolute value is equal to the number ρ_i , where $i = 1, \dots, s + 1$, and let $k^* = k_{s+1}$. We suppose that, if $s > 0$, then

$$(5) \quad e_i > 0$$

for $i = 1 \dots, s$. For given $y = y(\nu)$ in $\mathbb{C}^{m-1+\mathbb{N}}$, let

$$\omega_{n,y}(\nu) = \max(|y(\nu)|, \dots, |y(\nu + n - 1)|).$$

Then there exist $A > 0$, $m \in \mathbb{N}$, $\alpha^\wedge(\nu) > 0$ with $\nu \in m + \mathbb{N} - 1$ and the subspaces $V_{m,1}^\vee, \dots, V_{m,s+1}^\vee$ such that

$$\lim_{\nu \rightarrow \infty} \alpha^\wedge(\nu) = 0,$$

$$V_m = V_{m,1}^\vee \oplus \dots \oplus V_{m,s+1}^\vee, \dim_{\mathbb{C}}(V_{m,i}^\vee) = e_i, \quad 1 \leq i \leq s + 1,$$

and, if $y \in V_{m,\theta}^\vee$ for some $\theta \in \{1, \dots, s\}$, then

$$(6) \quad \exp(-A(\ln(\nu) + \nu^{1-1/k_\theta}))(\rho_\theta)^\nu \omega_n(y)(m) \leq \omega_{n,y}(\nu)$$

for $\nu \in m + \mathbb{N} - 1$; moreover, the spaces

$$V_{m,j}^\wedge = V_{m,j}^\vee \oplus \dots \oplus V_{m,s+1}^\vee,$$

where $j = 1 \dots, s + 1$, and, if $s \geq 1$ natural projections π_j ,

$$V_{m,j}^\wedge \mapsto V_{m,j}^\vee,$$

where $j = 1 \dots, s$, have the following properties:

if $y \in V_{m,\theta}^\wedge$ for some $\theta \in \{1, \dots, s\}$, then

$$(7) \quad \omega_{n,y}(\nu) \leq \exp(A(\ln(\nu) + \nu^{1-1/k_\theta}))(\rho_\theta)^\nu \omega_{n,y}(m),$$

$$(8) \quad (\omega_{n\pi_{\theta(y)}}(m) - \alpha(\nu)\omega_{n,y}(m))(\rho_{\theta})^{\nu} \\ \exp(-A(\ln(\nu) + \nu^{1-1/k_{\theta}})) \leq \omega_{n,y}(\nu),$$

where $\nu \in m + \mathbb{N} - 1$; if

$$(9) \quad k^* > 0,$$

and $y \in V_{m,s+1}^{\vee} (= V_{m,s+1}^{\wedge})$, then

$$(10) \quad |y(\nu)| \leq (A/\nu)^{\nu/k^*} \omega_{n,y}(m),$$

where $\nu \in m + \mathbb{N} - 1$.

Remark 1. It follows from the Theorem 7 that the space $V_{m,\theta}^{\wedge}$, where $\theta = 1, \dots, s+1$, does not depend from the construction and is defined uniquely by means of the equality

$$V_{m,\theta}^{\wedge} = \{y \in V_m : \limsup |y(\nu)|^{1/\nu} \leq \rho_{\theta}\}.$$

The presence of unknown $\alpha(\nu)$ (even tending to zero) in (8) constrict the possibilities of the application of this Theorem. Of course, in view of (6), in the case $e_{s+1} = 0$, $\theta = s$ this $\alpha(\nu)$ cannot play devil with us, because it vanishes then, but such happy case (see, for example, [75]-[76]) is rather an exception from the rule. One may attempt to estimate the specified value but it would be better to get rid from it at all. With this goal I prove here the following

Theorem 10. Let are fulfilled all the conditions of the Theorem 7. Let further V be an arbitrary linear subspace of V_m such that

$$V \cap V_{m,\theta+1} = \{0\},$$

where $\theta \in \{1 \dots, s\}$.

Then for this V there exists a constant $A^* = A^*(V) > 0$ such that

$$(11) \quad \exp(-A^*(\ln(\nu) + \nu^{1-1/k}))(\rho_{\theta})^{\nu} \omega_n(y)(m) \leq \omega_{n,y}(\nu)$$

where $y \in V$, $k = \max(k_1, \dots, k_s)$ and $\nu \in m + \mathbb{N} - 1$.

First I prove the following

Theorem 8. Let are fulfilled all the conditions of the Theorem 7. Let further

$$V_{m,j}^* = V_{m,1}^{\vee} \oplus \dots \oplus V_{m,j}^{\vee},$$

where $j = 1 \dots, s+1$, (and $V_{m,1}^* = V_{m,1}^{\vee}$.) Then vor $V = V_{m,\theta}^*$ with $\theta \in \{1, \dots, s\}$, holds the assertion of the Theorem 10.

Then I prove

Theorem 9. Let for some $\theta \in \{1 \dots, s\}$ is given a linear map ξ_{θ} of the space $V_{m,\theta}^*$ into $V_{m,\theta+1}^{\wedge}$. Let I_{θ}^* is the identity map $V_{m,\theta}^* \rightarrow V_{m,\theta}^*$ Then for

$$V = (I_{\theta}^* + \xi_{\theta})(V_{m,\theta}^*)$$

holds the assertion of the Theorem 10.

In the section 4 I prove the Theorem 10.

And in the section 5 I discuss, what will take place, if instead (1) the following conditions hold:

$$(12) \quad \lim_{\nu \rightarrow \infty} a_i(\nu) = a_i^\sim,$$

where $i = 0, \dots, n$,

$$(13) \quad a_n(\nu) = 1,$$

where $\nu \in \mathbb{N} - 1$.

§1. Some preparatory results.

Lemma 1. *Let $a \in \mathbb{N}$, $b \in \mathbb{N} - 1 + a$, $C > 0$ Then*

$$(14) \quad 0 < \sum_{\kappa=a}^b \ln(1 + C/\kappa) \leq$$

$$\ln(1 + C/a) + b \ln(1 + C/b) - a \ln(1 + C/a) + C \ln((b + C)/(a + C)).$$

Proof. See the proof of the Lemma 1 in [77]. ■

Corollary. *If $a \in \mathbb{N}$, $b \in \mathbb{N} + a - 1$, $b < 2a$, $C > 0$, then*

$$\sum_{\kappa=a}^b \ln(1 + C/\kappa) \leq 3C.$$

Proof. See the Proof of the Corollary of the Lemma 1 in [77]. ■

Lemma 2. *([51], Lemma 2, [44], Lemma 2, [72], Lemma 8.)*

Let $A \in \text{Mat}_n(\mathbb{C})$ and let k is a maximal order of its Jordan blocks. Then there exists a constant $\gamma^(A) > 0$ with the following properties:*

for any $\varepsilon > 0$ there exists a norm $p_{A,\varepsilon}$ on $\mathbb{C}^n$ such that

$$(15) \quad p_{A,\varepsilon} \leq \gamma^*(A)(\max(1, 1/\varepsilon)^{k-1}h,$$

$$(16) \quad h \leq \gamma^*(A)(\max(1, \varepsilon)^{k-1}p_{A,\varepsilon},$$

$$(17) \quad (p_{A,\varepsilon})^\sim \leq (\gamma^*(A))^2(\max(\varepsilon, 1/\varepsilon)^{k-1}h^\sim,$$

$$(18) \quad h^\sim \leq (\gamma^*(A))^2(\max(\varepsilon, 1/\varepsilon)^{k-1}(p_{A,\varepsilon})^\sim,$$

$$(19) \quad \|A\|_{sp} \leq (p_{A,\varepsilon})^\sim \leq \|A\|_{sp} + (\text{sign}(k - 1))\varepsilon,$$

where $\|A\|_{sp}$ denotes the maximum of the absolute values of eigenvalues of the matrix A . If, moreover,

$$(20) \quad \det(A) \neq 0, \quad \|A^{-1}\|_{sp}^{-1} > (\text{sign}(k - 1))\varepsilon,$$

then

$$(21) \quad \|A^{-1}\|_{sp} \leq$$

$$(p_{A,\varepsilon})^\sim(A^{-1}) \leq \left(\|A^{-1}\|_{sp}^{-1} - (\text{sign}(k-1))\varepsilon \right)^{-1} =$$

$$(\text{sign}(k-1)) \|A^{-1}\|_{sp} + \varepsilon \|A^{-1}\|_{sp} \left(\|A^{-1}\|_{sp}^{-1} - (\text{sign}(k-1))\varepsilon \right)^{-1}.$$

Proof. See the proof of the Lemma 8 in [72]. ■

Corollary. *If all the eigenvalues of the matrix A are simple, then*

$$(22) \quad (p_{A,\varepsilon})^\sim = \|A\|_{sp}.$$

If, moreover,

$$(23) \quad \det(A) \neq 0,$$

then

$$(24) \quad (p_{A,\varepsilon})^\sim(A^{-1}) = \left(\|A^{-1}\|_{sp} \right)^{-1}.$$

Proof. See the proof of the Corollary of the Lemma 8 in [72]. ■

Lemma 3. *Let all the conditions of the Theorem 7 are fulfilled, and let*

$$k = \max(k_1, \dots, k_s).$$

Then there exist $A > 0$, $m \in \mathbb{N}$ such that

$$(25) \quad \omega_{n,y}(\nu) \leq \exp(A(\ln(\nu) + \nu^{1-1/k}))(\rho_1)^\nu \omega_n(y)(m)$$

for any $y \in V_m$ and $\nu \in m + \mathbb{N} - 1$.

If, moreover, $k^* = 0$, then there exist $A > 0$, $m \in \mathbb{N}$ such that

$$(26) \quad \exp(-A(\ln(\nu) + \nu^{1-1/k}))(\rho_s)^\nu \omega_{n,y}(m) \leq \omega_{n,y}(\nu)$$

for any $y \in V_m$ and $\nu \in m + \mathbb{N} - 1$.

Proof. Since $V_m = V_{m,1}^\wedge$, it follows that the inequality (7) holds with $\theta = 1$ for any $y \in V_m$. For the full proof of the Lemma let us make some not large changes in the proof of the Lemma 2 in [77].

The condition $k^* = 0$ implies the inequality

$$(27) \quad a_0^\sim = T(0) \neq 0,$$

and for p in Theorem 6 of the paper [72] the equality $p = n$. Let

$$(28) \quad A(\nu) = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & 1 \\ -a_0(\nu) & -a_1(\nu) & -a_2(\nu) & \dots & -a_{n-1}(\nu) \end{pmatrix},$$

where $\nu \in \mathbb{N} - 1$, and let

$$(29) \quad A^\sim = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & 1 \\ -a_0^\sim & -a_1^\sim & -a_2^\sim & \dots & -a_{n-1}^\sim \end{pmatrix}$$

Let λ_j , where $j = 1, \dots, r$, is the sequence of all the mutually distinct roots of the polynomial (4) and k_j^* is the multiplicity of the root λ_j . Then

$$k = \sup\{k_j^* : j = 1, \dots, r\}.$$

Clearly,

$$\sum_{j=0}^r k_j^* = n,$$

$$\rho_s \leq |\lambda_j| \leq \rho_1 = \|A\|_{sp} < R = h^\sim(A^\sim) + 1,$$

where $j = 1, \dots, r$. In view of (1), there exists $C_1 > 0$ such that

$$(30) \quad h^\sim(A(\nu) - A^\sim) \leq C_1/(\nu + 1).$$

Therefore, according to the Lemma 2, for any $\varepsilon > 0$ the following inequalities holds

$$(31) \quad p_{A^\sim, \varepsilon}(A(\nu) - A^\sim) \leq (\gamma^*(A^\sim))^2 (\max(\varepsilon, 1/\varepsilon))^{k-1} h^\sim(A(\nu) - A^\sim) \leq (\gamma^*(A^\sim))^2 (\max(\varepsilon, 1/\varepsilon))^{k-1} C_1/(\nu + 1),$$

$$(32) \quad p_{A^\sim, \varepsilon}(A(\nu)) \leq \rho_1 + \text{sign}(k - 1)\varepsilon + (\gamma^*(A^\sim))^2 (\max(\varepsilon, 1/\varepsilon))^{k-1} C_1/(\nu + 1) \leq (\rho_1 + (\text{sign}(k - 1))\varepsilon) \left(1 + (\gamma^*(A^\sim))^2 (\max(\varepsilon, 1/\varepsilon))^{k-1} \frac{C_1}{\rho_1(\nu + 1)} \right),$$

$$(33) \quad \ln(p_{A^\sim, \varepsilon}(A(\nu))) \leq \ln(\rho_1) + \ln(1 + (\text{sign}(k - 1))C_{1,1}\varepsilon) + \ln(1 + (\max(\varepsilon, 1/\varepsilon))^{k-1} C_{1,2}/(\nu + 1)),$$

where

$$C_{1,1} = 1/\rho_1, \quad C_{1,2} = (\gamma^*(A^\sim))^2 C_1/\rho_1$$

and $\nu \in \mathbb{N}$.

We consider first the case $k > 1$. For given $\nu \in \mathbb{N} + 1$ we take $d \in \mathbb{N}$ and $a_i \in [1, \nu] \cap \mathbb{N}$, where $i = 0, \dots, d$, in such a way that

$$(34) \quad a_0 = 1, \quad a_d = \nu, \quad a_{i-1} < a_i \leq 2a_{i-1},$$

where $i = 1, \dots, d$, and

$$(35) \quad d \leq \frac{\ln(\kappa)}{\ln(2)} + 1.$$

According to the Corollary of the Lemma 1 and (34) – (35),

$$(36) \quad \sum_{\kappa=a_{i-1}}^{a_i-1} \ln(p_{A^\sim, \varepsilon}(A(\kappa))) \leq$$

$$(a_i - a_{i-1}) \ln(\rho_1) + (a_i - a_{i-1}) \ln(1 + C_{1,1}\varepsilon) +$$

$$\sum_{\kappa=a_{i-1}}^{a_i-1} \ln(1 + C_{1,2}(\max(\varepsilon, 1/\varepsilon))^{k-1}/(\kappa + 1)) \leq$$

$$(a_i - a_{i-1})(\ln(\rho_1) + \ln(1 + \varepsilon/\rho_1)) + 3C_{1,2}(\max(\varepsilon, 1/\varepsilon))^{k-1},$$

where $i = 1, \dots, d$. We take now in (36) $\varepsilon = \varepsilon_i = (a_{i-1})^{-1/k}$. Then we obtain the inequality

$$\sum_{\kappa=a_{i-1}}^{a_i-1} \ln(p_{A^\sim, \varepsilon_i}(A(\kappa))) \leq$$

$$(a_i - a_{i-1}) \ln(\rho_1) + (C_{1,1} + 3C_{1,2})(a_{i-1})^{1-1/k} =$$

$$(a_i - a_{i-1}) \ln(\rho_1) + O(2^{(i-1)(1-1/k)}),$$

where $i = 1, \dots, d$. Therefore

$$(37) \quad \ln(p_{A^\sim, \varepsilon_i} \left(\prod_{\kappa=1}^{a_i-a_{i-1}} A(a_i - \kappa) \right)) \leq$$

$$(a_i - a_{i-1}) \ln(\rho_1) + O(2^{(i-1)(1-1/k)})$$

and, in view of (18),

$$(38) \quad \ln(h^\sim \left(\prod_{\kappa=1}^{a_i-a_{i-1}} A(a_i - \kappa) \right)) \leq$$

$$\ln((\gamma^*(A^\sim))^2)(\max(\varepsilon, 1/\varepsilon))^{k-1} + (a_i - a_{i-1}) \ln(\rho_1) + O(2^{(i-1)(1-1/k)}) \leq$$

$$2 \ln(\gamma^*(A^\sim)) + (i-1)(1-1/k) \ln(2) + (a_i - a_{i-1}) \ln(\rho_1) + O(2^{(i-1)(1-1/k)}) =$$

$$(i-1)(1-1/k) \ln(2) + (a_i - a_{i-1}) \ln(\rho_1) + O(2^{(i-1)(1-1/k)}),$$

where $i = 1, \dots, d$. Consequently,

$$(39) \quad \ln \left(h^\sim \left(\prod_{\kappa=1}^{nu-1} A(\nu - \kappa) \right) \right) =$$

$$\ln \left(h^\sim \left(\left(\prod_{i=1}^d \prod_{\kappa=1}^{a_i-a_{i-1}} A(a_i - \kappa) \right) \right) \right) =$$

$$= \nu \ln(\rho_1) + O(\nu^{(1-1/k)}),$$

where $\nu \in \mathbb{N}$.

If $k = 1$, then, according to (33),

$$(40) \quad \begin{aligned} \ln(p_{A^\sim,1}^\sim(A(\nu))) &\leq \ln(\rho_1) + C_{1,2}/(\nu + 1), \\ \ln \left(h^\sim \left(\prod_{\kappa=1}^{\nu-1} A(\nu - \kappa) \right) \right) &\leq \\ \ln \left((\gamma^*(A^\sim))^2 p_{A^\sim,1}^\sim \left(\prod_{\kappa=1}^{\nu-1} A(\nu - \kappa) \right) \right) &\leq \\ \nu \ln(\rho_1) + O(\ln(e\nu)). \end{aligned}$$

where $\nu \in \mathbb{N}$. In view of (39) – (40),

$$(41) \quad \begin{aligned} \ln \left(h^\sim \left(\prod_{\kappa=1}^{\nu-1} A(\nu - \kappa) \right) \right) &= \\ = \nu \ln(\rho_1) + O((\nu + 1)^{(1-1/k)}) + O(\ln(e\nu)), \end{aligned}$$

where $\nu \in \mathbb{N}$.

As in Section 3 of [72], let K denotes one of the fields $\mathbb{R}$ or $\mathbb{C}$ and L denotes a linear normed space over K with norm $p = p(x)$. If $L = K^n$, we fix as $p = p(x)$, where $x \in K^n$, the maximum of the absolute values of coordinates of the element x in the standard basis, i.e.

$$(42) \quad p(x) = h(x) = \sup(\{|x_1|, \dots, |x_n|\}),$$

where

$$x = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}.$$

If L is a Banach space with the norm p , then K -algebra of all the linear continuous operators acting in L will be denoted by $\mathfrak{M}^\wedge(L)$, and the norm on $\mathfrak{M}^\wedge(L)$, associated with the norm p will be denoted by $p^\sim$. So,

$$p^\sim(A) = \sup(\{p(AX) : X \in L, p(X) \leq 1\}).$$

It is well known that the associated with h norm on $Mat_n(\mathbb{C})$ is defined as follows

$$(43) \quad h^\sim(A) = \sup \left(\left\{ \sum_{k=1}^n |a_{i,j}| : i = 1, \dots, n \right\} \right),$$

where $A = (a_{i,k}) \in Mat_n(\mathbb{C})$. The norms h and $h^\sim$ coincide respectively with the norms q_∞ and $q_\infty^\sim$ considered in section 6 of the paper [68].

Let $m \in \mathbb{N}$, and let $E_m(L)$ be the set $L^{m-1+\mathbb{N}}$ of all the maps of the set $m-1+\mathbb{N}$ into L . The set $E_m(L)$ is a linear space over K , where the multiplication of the elements by the number from K and addition of the elements are defined coordinate-wise. The subspace of $E_m(L)$ composed by

all the constant maps is isomorphic to the space L , and we identify this subspace with L .

As in Section 4 of [72], for any $y \in E_m(\mathbb{C})$ and $n \in \mathbb{N}$ let $Y_{n,y}$ and $Y_{n,y}^\#$ denote the elements in the space $E_m(\mathbb{C}^n)$, which are determined respectively by means the following equalities:

$$(44) \quad Y_{n,y}(\nu) = \begin{pmatrix} y(\nu) \\ \vdots \\ y(\nu + n - 1) \end{pmatrix},$$

$$(45) \quad Y_{n,y}^\#(\nu) = \begin{pmatrix} 0 \\ \vdots \\ 0 \\ y(\nu) \end{pmatrix},$$

where $\nu \in m - 1 + \mathbb{N}$. Let is fixed $m \in \mathbb{N}$. If $y = y(\nu)$ is a solution of the equation (3) for $\nu \in \mathbb{N} + m - 1$, then

$$Y_{n,y}(\nu) = \left(\prod_{\kappa=1}^{\nu-m} A(\nu - \kappa) \right) Y_{n,y}(m),$$

where $\nu \in \mathbb{N}$, and, in view of (41),

$$(46) \quad \begin{aligned} \omega_{n,y}(\nu) &= h(Y_{n,y}(\nu)) \leq \\ &h^\sim \left(\prod_{\kappa=1}^{\nu-m} A(\nu - \kappa) \right) h(Y_{n,y}(m)) = \\ &\exp(O(1)(\nu^{1-1/k} + \ln(\nu))) (\rho_1)^\nu \omega_{n,y}(m), \end{aligned}$$

where $\nu \in \mathbb{N} + m - 1$. So, with $m = 1$ the asserted by the Lemma the upper estimate (25) of the value $\omega_{n,y}(\nu) (\rho_\theta)^{-\nu} = \omega_{n,y}(\nu) (\rho_1)^{-\nu}$ is obtained. We shall take now

$$(47) \quad \varepsilon \in (0, \rho_s/2).$$

Then, in view of (21),

$$(48) \quad \begin{aligned} 1/\rho_s &\leq (p_{A,\varepsilon})^\sim(A^{-1}) \leq \\ 1/\rho_s + 2(\text{sign}(k-1))\varepsilon/\rho_s^2 &\leq 2/\rho_s. \end{aligned}$$

Let is fixed

$$(49) \quad m \in \max([(2/\rho_s)^k], [2C_1 h^\sim((A^\sim)^{-1})]) + \mathbb{N}.$$

Then

$$(50) \quad \begin{aligned} h^\sim((A^\sim)^{-1}(A(\nu) - A^\sim)) &\leq h^\sim((A^\sim)^{-1}) C_1/(\nu + 1) \leq \\ h^\sim((A^\sim)^{-1}) C_1/([2C_1 h^\sim((A^\sim)^{-1})] + 1) &\leq 1/2, \end{aligned}$$

if $\nu \in m - 1 + \mathbb{N}$, the matrix $E + ((A^\sim)^{-1}(A(\nu) - A^\sim))$ is invertible,

$$h^\sim \left((E + ((A^\sim)^{-1}(A(\nu) - A^\sim)))^{-1} \right) \leq 2,$$

if $\nu \in m - 1 + \mathbb{N}$, there exists the matrix

$$(A(\nu))^{-1} = (E + ((A^\sim)^{-1}(A(\nu) - A^\sim)))^{-1} (A^\sim)^{-1},$$

if $\nu \in m - 1 + \mathbb{N}$, moreover,

$$\begin{aligned} h^\sim ((A(\nu))^{-1}) &\leq h^\sim \left((E + ((A^\sim)^{-1}(A(\nu) - A^\sim)))^{-1} \right) h^\sim ((A^\sim)^{-1}) \leq \\ &2h^\sim ((A^\sim)^{-1}) \end{aligned}$$

and, finally,

$$\begin{aligned} h^\sim ((A(\nu))^{-1} - (A^\sim)^{-1}) &= \\ h^\sim ((A(\nu))^{-1} (A^\sim - A(\nu)) (A^\sim)^{-1}) &\leq \\ h^\sim ((A(\nu))^{-1}) h^\sim ((A^\sim - A(\nu))) h^\sim ((A^\sim)^{-1}) &\leq C_2/(\nu + 1), \end{aligned}$$

where

$$C_2 = (h^\sim ((A^\sim)^{-1}))^2 C_1$$

and $\nu \in m - 1 + \mathbb{N}$. Therefore

$$\begin{aligned} (51) \quad p_{A^\sim, \varepsilon}((A(\nu))^{-1} - (A^\sim)^{-1}) &\leq \\ (\gamma^*(A^\sim))^2 (\max(\varepsilon, 1/\varepsilon))^{k-1} h^\sim((A(\nu))^{-1} - (A^\sim)^{-1}) &\leq \\ (\gamma^*(A^\sim))^2 (\max(\varepsilon, 1/\varepsilon))^{k-1} C_2/(\nu + 1), \end{aligned}$$

where $\nu \in m - 1 + \mathbb{N}$. In view of (51) and (21),

$$\begin{aligned} (52) \quad p_{A^\sim, \varepsilon}((A(\nu))^{-1}) &\leq 1/\rho_s + \\ 2(\text{sign}(k-1))\varepsilon/\rho_s^2 &\leq 2/\rho_s + (\gamma^*(A^\sim))^2 (\max(\varepsilon, 1/\varepsilon))^{k-1} C_2/(\nu + 1) \leq \\ (1/\rho_s + 2(\text{sign}(k-1))\varepsilon/\rho_s^2) &(1 + (\gamma^*(A^\sim))^2 (\max(\varepsilon, 1/\varepsilon))^{k-1} \rho_s C_2 \rho_s/(\nu + 1)), \end{aligned}$$

$$\begin{aligned} (53) \quad \ln(p_{A^\sim, \varepsilon}((A(\nu))^{-1})) &\leq \ln(1/\rho_s) + \\ \ln(1 + (\text{sign}(k-1))C_{2,1}\varepsilon) &+ \ln(1 + (\max(\varepsilon, 1/\varepsilon))^{k-1} C_{2,2}/(\nu + 1)), \end{aligned}$$

where

$$C_{2,1} = 2/\rho_s, \quad C_{2,2} = (\gamma^*(A^\sim))^2 C_2 \rho_s$$

and $\nu \in \mathbb{N} - 1 + m$. We take $\nu \in m - 1 + \mathbb{N}$.

We consider first the case $k > 1$ again now. For given $\nu \in \mathbb{N} + m$ we take $d \in \mathbb{N}$ and $a_i \in [m, \nu] \cap \mathbb{N}$, where $i = 0, \dots, d$, in such a way that

$$(54) \quad a_0 = m, \quad a_d = \nu, \quad a_{i-1} < a_i \leq 2a_{i-1} \leq m2^i,$$

where $i = 1, \dots, d$, and

$$(55) \quad d \leq \frac{\ln(\nu)}{\ln(2)} + 1.$$

According to the Corollary of the Lemma 1 and (54) – (55),

$$\begin{aligned}
 (56) \quad & \sum_{\kappa=a_{i-1}}^{a_i-1} \ln(p_{A^\sim, \varepsilon}((A(\kappa))^{-1})) \leq \\
 & (a_i - a_{i-1}) \ln(1/\rho_s) + (a_i - a_{i-1}) \ln(1 + C_{2,1}\varepsilon) + \\
 & \sum_{\kappa=a_{i-1}}^{a_i-1} \ln(1 + (\max(\varepsilon, 1/\varepsilon))^{k-1} C_{2,2}/(\kappa + 1)) \leq \\
 & (a_i - a_{i-1})(\ln(1/\rho_s) + \ln(1 + C_{2,1}\varepsilon)) + 3(\gamma^*(A^\sim))^2(\max(\varepsilon, 1/\varepsilon))^{k-1} C_{2,2},
 \end{aligned}$$

where $i = 1, \dots, d$. We take now in (56) $\varepsilon = \varepsilon_i = (a_{i-1})^{-1/k}$. Then, in view of (49),

$$a_{i-1} \geq m > (2/\rho_s)^k, (a_{i-1})^{1/k} > 2/\rho_s, \varepsilon_i < \rho_s/2,$$

where $i = 1, \dots, d$, and therefore (47) and (56) hold. Consequently,

$$\begin{aligned}
 (57) \quad & \sum_{\kappa=a_{i-1}}^{a_i-1} \ln(p_{A^\sim, \varepsilon}((A(\kappa))^{-1})) \leq \\
 & (a_i - a_{i-1}) \ln(1/\rho_s) + (C_{2,1} + 3C_{2,2})(a_{i-1})^{1-1/k} = \\
 & (a_i - a_{i-1}) \ln(1/\rho_s) + O(2^{(i-1)(1-1/k)}),
 \end{aligned}$$

where $i = 1, \dots, d$. Therefore

$$\begin{aligned}
 (58) \quad & \ln(p_{A^\sim, \varepsilon_i} \left(\prod_{\kappa=a_{i-1}}^{a_i-1} (A(\kappa))^{-1} \right)) \leq \\
 & (a_i - a_{i-1}) \ln(\rho_1) + O(2^{(i-1)(1-1/k)})
 \end{aligned}$$

and, in view of (18),

$$\begin{aligned}
 (59) \quad & \ln(h^\sim \left(\prod_{\kappa=1}^{a_i-a_{i-1}} (A(\kappa))^{-1} \right)) \leq \\
 & \ln((\gamma^*(A^\sim))^2)(\max(\varepsilon_i, 1/\varepsilon_i))^{k-1} + (a_i - a_{i-1}) \ln(\rho_1) + O(2^{(i-1)(1-1/k)}) \leq \\
 & 2 \ln(\gamma^*(A^\sim)) + (i-1)(1-1/k) \ln(2) + (a_i - a_{i-1}) \ln(\rho_1) + O(2^{(i-1)(1-1/k)}) = \\
 & (i-1)(1-1/k) \ln(2) + (a_i - a_{i-1}) \ln(\rho_1) + O(2^{(i-1)(1-1/k)}),
 \end{aligned}$$

where $i = 1, \dots, d$. Consequently,

$$\begin{aligned}
 (60) \quad & \ln \left(h^\sim \left(\prod_{\kappa=m}^{\nu-1} (A(\kappa))^{-1} \right) \right) = \\
 & \ln \left(h^\sim \left(\left(\prod_{i=1}^d \prod_{\kappa=1}^{a_i-a_{i-1}} (A(\kappa))^{-1} \right) \right) \right) = \\
 & = \nu \ln(1/\rho_s) + O(\nu^{(1-1/k)}),
 \end{aligned}$$

where $\nu \in \mathbb{N} + m$.

If $k = 1$, then, according to (53),

$$(61) \quad \begin{aligned} \ln(p_{A^\sim,1}(A(\nu))) &\leq \ln(1/\rho_s) + C_{2,2}/(\nu + 1), \\ \ln \left(h^\sim \left(\prod_{\kappa=m}^{\nu-1} (A(\kappa))^{-1} \right) \right) &\leq \\ \ln \left((\gamma^*(A^\sim))^2 p_{A^\sim,1} \left(\prod_{\kappa=m}^{\nu-1} (A(\kappa))^{-1} \right) \right) &\leq \\ \nu \ln(1/\rho_s) + O(\ln(e\nu)). \end{aligned}$$

where $\nu \in \mathbb{N}$. In view of (60) – (61),

$$(62) \quad \begin{aligned} \ln \left(h^\sim \left(\prod_{\kappa=m}^{\nu-1} A(\nu - \kappa) \right) \right) &= \\ = \nu \ln(1/\rho_s) + O(\nu^{1-1/k}) + O(\ln(e\nu)), \end{aligned}$$

where $\nu \in \mathbb{N} + m$. Since

$$\begin{aligned} Y_{n,y}(m) &= \left(\prod_{\kappa=1}^{\nu-m} A(\nu - \kappa) \right)^{-1} Y_{n,y}(\nu) = \\ &= \left(\prod_{\kappa=m}^{\nu-1} (A(\kappa))^{-1} \right) Y_{n,y}(\nu), \end{aligned}$$

it follows that

$$(63) \quad \begin{aligned} \omega_{n,y}(m) &= h(Y_{n,y}(m)) \leq \\ h^\sim \left(\prod_{\kappa=m}^{\nu-1} (A(\kappa))^{-1} \right) h(Y_{n,y}(\nu)) &= \\ \exp(O(1)(\nu^{1-1/k} + \ln(\nu))) (\rho_1)^{-\nu} \omega_{n,y}(\nu), \end{aligned}$$

where $\nu \in \mathbb{N} + m$. So, with m from (49) the asserted by the Lemma the lower estimate of the value $\omega_{n,y}(\nu) (\rho_s)^{-\nu}$ is obtained. ■

Remark 1. In the case $s = 1$, $k^* = 0$ the assertions of the Lemma and Theorem 7 coincide.

Lemma 4 (Perron's decomposition Lemma, [22], Hilfsatz 3)). Let the characteristic polynomial (4) is represented as product

$$(64) \quad T(z) = T_1(z)T_2(z),$$

where

$$(65) \quad T_1(z) = \sum_{\alpha=0}^p b_\alpha^\sim z^\alpha, \quad T_2(z) = \sum_{\beta=0}^q u_\beta^\sim z^\beta,$$

with $b_p^\sim = u_q^\sim = a_n^\sim = 1$ and absolute value of each root of $T_1(z)$ is greater than the absolute value of each root of $T_2(z)$.

Then there exist $m \in \mathbb{N}$, $b_\alpha(\nu) \in \mathbb{C}$, $\alpha = 0, \dots, p$, $\nu \in \mathbb{N} + m - 1$, and $u_\beta(\nu) \in \mathbb{C}$, $\beta = 0, \dots, q$, $\nu \in \mathbb{N} + m - 1$, such that

$$(66) \quad \lim_{\nu \rightarrow \infty} b_\alpha(\nu) = b_\alpha^\sim, \alpha = 0, \dots, p, b_p(\nu) = 1, b_0(\nu) \neq 0,$$

$$(67) \quad \lim_{\nu \rightarrow \infty} u_\beta(\nu) = u_\beta^\sim, \beta = 0, \dots, q, u_q(\nu) = 1,$$

where $\nu \in \mathbb{N} + m - 1$, and, moreover, the equation (3) is equivalent to the equation

$$\sum_{\alpha=0}^p b_\alpha(\nu) y(\nu + \alpha) = r(\nu),$$

where $\nu \in \mathbb{N} - 1 + m$ and $r(\nu)$ satisfies to the equation

$$\sum_{\beta=0}^q u_\beta(\nu) r(\nu + \beta) = 0$$

with $\nu \in \mathbb{N} - 1 + m$.

Lemma 5. Let all the conditions of the Perron's decomposition Lemma are fulfilled and $a_k(\nu) - a_k^\sim = O(1/(\nu + 1))$, $k = 0, \dots, n$, when $\nu \rightarrow \infty$.

Then for $b_\alpha(\nu)$ with $\alpha = 0, \dots, p$, and $u_\beta(\nu)$ with $\beta = 0, \dots, q$, from the assertion of the Perron's decomposition Lemma the following conditions are fulfilled (cf. with (66) and (67)):

$$b_\alpha(\nu) - b_\alpha^\sim = O(1/\nu), \alpha = 0, \dots, p, b_0(\nu) = 1, b_q(\nu) \neq 0,$$

$$u_\beta(\nu) - u_\beta^\sim = O(1/\nu), \beta = 0, \dots, q, u_0(\nu) = 1,$$

where $\nu \in \mathbb{N} - 1 + m$.

Proof. The Lemma is direct corollary of the Theorem 5 in [68]. ■

§2. Proof of the theorem 8.

We use below the notations of the section 3 in [77]. Let K be one of the fields $\mathbb{R}$ or $\mathbb{C}$. Let L be a linear normed space over K with a norm $p(x)$. In the case $L = K^n$ we fix as $p(x)$, where $x \in K^n$, the maximum of the absolute values of coordinates of x in the standard basis, i.e.

$$(68) \quad p(x) = h(x) = \sup(\{|x_1|, \dots, |x_n|\}),$$

where

$$x = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}.$$

If L is a Banach space with the norm p , then K -algebra of all the linear continuous operators acting in L will be denoted by $\mathfrak{M}^\wedge(L)$, and the norm on $\mathfrak{M}^\wedge(L)$, associated with the norm p will be denoted by $p^\sim$. So,

$$p^\sim(A) = \sup(\{p(AX) : X \in L, p(X) \leq 1\}).$$

It is well known that the associated with h norm on $Mat_n(\mathbb{C})$ is defined as follows

$$(69) \quad h^\sim(A) = \sup \left(\left\{ \sum_{k=1}^n |a_{i,j}| : i = 1, \dots, n \right\} \right),$$

where $A = (a_{i,k}) \in Mat_n(\mathbb{C})$. The norms h and $h^\sim$ coincide respectively with the norms q_∞ and $q_\infty^\sim$ considered in [68], section 6. Let $m \in \mathbb{N} - 1$, and we denote by $E_m(L)$ the set $L^{m-1+\mathbb{N}}$ of all the maps of the set $m-1+\mathbb{N}$ into L . The set $E_m(L)$ is a linear space over K , where the multiplication of the elements by the number from K and addition of the elements is defined coordinate-wise. The subspace of $E_m(L)$ composed by all the constant maps is isomorphic to L , and we identify this subspace with L .

We denote by $\mathfrak{M}^\vee(L)$ the space of all the K -linear maps of the space L in L . If $\phi \in \mathfrak{M}^\vee(L)$ and $\psi \in \mathfrak{M}^\vee(L)$, then $\phi \circ \psi$ denotes the composition of operators ϕ and ψ , so that $(\phi \circ \psi)f = \phi(\psi f)$ for each $f \in L$. For $x \in E_m(L)$ let

$$p_{m,\infty}(x) = \sup(\{p(x(\nu)) : \nu \in m-1+\mathbb{N}\}).$$

Let further

$$E_{m,\infty}(L) = \{x \in E_m(L) : p_{m,\infty}(x) \neq \infty\},$$

$$E_{m,0}(L) = \{x \in E_m(L) : \lim_{\nu \rightarrow \infty} p(x(\nu)) = 0\},$$

$$E_m^\rightarrow(L) = L + E_{m,0}(L).$$

Clearly, the space $E_m^\rightarrow(L)$ consists of all the $y \in E_m(L)$, for which there exists

$$\lim(y) = \lim_{\nu \rightarrow \infty} (y(\nu)).$$

Let $m \in \mathbb{N} - 1$, $\mu \in m-1+\mathbb{N}$ and let $r_{m,\mu}$ be the operator of restriction of the elements $y \in E_m(L)$ on the set $m-1+\mathbb{N}$. Clearly, the map $r_{m,\mu}$ is an epimorphism of the space $E_m(L)$ onto the space $E_\mu(L)$. If L is a K -algebra, then $E_m(L)$ is a K -algebra, where the multiplication and addition of the elements is defined coordinate-wise; so, in this case $r_{m,\mu}$ is an epimorphism of K -algebra $E_m(L)$ onto K -algebra $E_\mu(L)$. If L be an algebra with unity, let L^* denotes the group of all its invertible elements. Then

$$(L^*)^{m-1+\mathbb{N}} \subset L^{m-1+\mathbb{N}};$$

we denote below $(L^*)^{m-1+\mathbb{N}}$ by $E_m(L^*)$. Clearly,

$$E_m(L^*) = (E_m(L))^*.$$

Let $L = \mathbb{C}^n$, $y \in E_m(L)$, and let $y_i(\nu)$ denotes the i -th coordinate of the element $y(\nu)$, where $i = 1, \dots, n$, $\nu \in m-1+\mathbb{N}$; then the space $(E_m(\mathbb{C}))^n$ contains an element $\omega(y)$, which has $y_i(\nu)$ as the value of its i -th coordinate at the point $\nu \in m-1+\mathbb{N}$. So we obtain the natural isomorphism ω of the algebra $E_m(\mathbb{C}^n)$ onto $(E_m(\mathbb{C}))^n$. This map ω induces an isomorphism of the algebra $E_m(Mat_n(\mathbb{C}))$ onto $Mat_n(E_m(\mathbb{C}))$. If L is a K -algebra, then each element $a \in E_m(L)$ determines an acting on the space $E_m(L)$ K -linear operator $\mu_a \in \mathfrak{M}^\vee(E_m(L))$, which turns any $y \in E_m(L)$ into $\mu_a y = ay$. On

the space $E_m(L)$ acts also K -linear operator $\nabla \in \mathfrak{M}^\vee(L)$, which turns any element $y \in E_m(L)$ in the $\nabla y \in E_m(L)$ such that

$$(\nabla y)(\nu) = y(\nu + 1)$$

for any $\nu \in m - 1 + \mathbb{N}$. Let us consider the subring $\mathfrak{A}_m(L)$ of the ring $\mathfrak{M}^\vee(\mathfrak{L})$ generated by the operator ∇ and by all the operators μ_a , where $a \in E_m(L)$. Clearly,

$$(70) \quad \mu_a \circ \nabla^r \circ \mu_b \circ \nabla^s = \mu_{a \nabla^r b_k} \circ \nabla^{r+s},$$

where $\{r, s\} \subset \mathbb{N} - 1$, $\{a, b\} \subset E_m(L)$. For each $\alpha \in \mathfrak{A}_m(L) \setminus \{0_{\mathfrak{A}_m(L)}\}$ are uniquely defined the number $\deg(\alpha)$ and representation of α in the form

$$(71) \quad \alpha = \sum_{k=0}^{\deg(\alpha)} \mu_{a_k} \circ \nabla^k,$$

where $a_k \in E_m(L)$ for $k = 0, \dots, \deg(\alpha)$ and $a_{\deg(\alpha)} \neq 0_{E_m(L)}$. Clearly, (71) may be rewritten in the form

$$(72) \quad \alpha = \sum_{k=0}^{\infty} \mu_{a_k} \circ \nabla^k,$$

where $a_k = 0_{E_m(L)}$ for $k \in \deg(\alpha) + \mathbb{N}$. It follows from (70) that $\mathfrak{A}_m(L)$ is a graduated algebra, and if

$$(73) \quad \beta = \sum_{r=0}^p \mu_{b_r} \circ \nabla^r \in \mathfrak{A}_m(L),$$

$$(74) \quad \gamma = \sum_{s=0}^q \mu_{c_s} \circ \nabla^s \in \mathfrak{A}_m(L),$$

then

$$(75) \quad \beta\gamma = \sum_{k=0}^{p+q} \sum_{\substack{\leq r \leq p \\ 0 \leq s \leq q \\ r+s=k}} \mu_{b_r \nabla^r c_s} \circ \nabla^{r+s};$$

clearly, $\deg(\beta\gamma) = \deg(\beta) + \deg(\gamma)$, if $b_p(\nu)^r c_q(\nu + p)$ is different from 0 at least for one $\nu \in m - 1 + \mathbb{N}$. Let $\mathfrak{A}_m^\rightarrow(L)$ be the ring generated by the operator ∇ and by all the operators μ_a , where $a \in E_m^\rightarrow(L)$. Since $\nabla a \in E_m^\rightarrow(L)$, if $a \in E_m^\rightarrow(L)$, it follows, in view of (70), that $\mathfrak{A}_m^\rightarrow(L)$ is a graduated subalgebra $\mathfrak{A}_m^\rightarrow(L)$ of the algebra $\mathfrak{A}_m(L)$, each $\alpha \in \mathfrak{A}_m(L) \setminus \{0_{\mathfrak{A}_m(L)}\}$ admits a representation in the form (71) with $a_k \in E_m^\rightarrow(L)$ for $k = 0, \dots, \deg(\alpha)$ and $a_{\deg(\alpha)} \neq 0_{E_m(L)}$; to each such α corresponds the limit operator

$$(76) \quad \lim(\alpha) = \sum_{k=0}^{\deg(\alpha)} \mu_{\lim(a_k)} \circ \nabla^k,$$

and polynomial

$$(77) \quad P(\alpha, z) = \sum_{k=0}^{\deg(\alpha)} \lim(a_k) z^k \in L[z].$$

If $\alpha = 0_{\mathfrak{A}_m(L)}$, then we put

$$\lim(\alpha) = 0_{\mathfrak{A}_m(L)}, P(\alpha, z) = 0_{L[z]}.$$

The equality (70) shows that the map

$$(78) \quad \alpha \rightarrow P(\alpha, z)$$

is an epimorphism of the algebra $\mathfrak{A}_m^{\rightarrow}(L)$ on the algebra $L[z]$ (if the algebra L is noncommutative, then we can treat the algebra $L[z]$ as a semigroup ring of the semigroup $(\mathbb{N} - 1, +)$ over the algebra L). We note that, if $\alpha \in \mathfrak{A}_m(\mathbb{C})$, then, clearly, $\text{Ker}(\alpha)$ coincides with the linear space of all the solutions of the equation (3), and, moreover, if $\alpha \in \mathfrak{A}_m^{\rightarrow}(\mathbb{C})$, then the corresponding to α equation (3) is an equation of the Poincaré type and $P(\alpha, z)$ is its characteristical polynomial.

Let L be an algebra with unity. The set of all $\alpha \in \mathfrak{A}_m(L) \setminus \{0_{\mathfrak{A}_m(L)}\}$, which have the representation (71) with $a_{\deg(\alpha)} \in E_m(L^*)$ will be denoted further by $\mathfrak{A}_m(L)^{\circ}$. The set of all the elements $\alpha \in \mathfrak{A}_m(L) \setminus \{0_{\mathfrak{A}_m(L)}\}$, which have the representation (71) with $a_{\deg(\alpha)} = 1_{E_m(L)}$ will be denoted further by $\mathfrak{A}_m(L)^{\vee}$. The set of all the $\alpha \in \mathfrak{A}_m(L) \setminus \{0_{\mathfrak{A}_m(L)}\}$, which have the representation (71) with $a_0 \in E_m(L^*)$, will be denoted further by $\mathfrak{A}_m(L)^{\wedge}$. Let further

$$\mathfrak{A}_m(L)^{\circ\wedge} = \mathfrak{A}_m(L)^{\circ} \bigcap \mathfrak{A}_m(L)^{\wedge}, \mathfrak{A}_m(L)^{\vee\wedge} = \mathfrak{A}_m(L)^{\vee} \bigcap \mathfrak{A}_m(L)^{\wedge},$$

$$\mathfrak{A}_m(L)^{\circ\rightarrow} = \mathfrak{A}_m(L)^{\circ} \bigcap \mathfrak{A}_m(L)^{\rightarrow}, \mathfrak{A}_m(L)^{\vee\rightarrow} = \mathfrak{A}_m(L)^{\vee} \bigcap \mathfrak{A}_m(L)^{\rightarrow},$$

$$\mathfrak{A}_m(L)^{\circ\wedge\rightarrow} = \mathfrak{A}_m(L)^{\circ\wedge} \bigcap \mathfrak{A}_m(L)^{\rightarrow}, \mathfrak{A}_m(L)^{\vee\wedge\rightarrow} = \mathfrak{A}_m(L)^{\vee\wedge} \bigcap \mathfrak{A}_m(L)^{\rightarrow},$$

Clearly, $\mathfrak{A}_m(L)^{\circ}$ consists of epimorphisms of the space $E_m(L)$ onto $E_m(L)$. The above map $r_{m,\mu}$ induces epimorphism $r_{m,\mu}^{\triangleright}$ of the algebra $\mathfrak{A}_m(L)$ on the algebra $\mathfrak{A}_{\mu}(L)$ defined as follows:

if $\alpha \in \mathfrak{A}_m(L)$,

$$(79) \quad \alpha = \sum_{k=0}^n \mu_{a_k} \circ \nabla^k,$$

then

$$(80) \quad r_{m,\mu}^{\triangleright}(\alpha) = \sum_{k=0}^n \mu_{r_{m,\mu}(a_k)} \circ \nabla^k,$$

where the operator ∇ in (79) acts in $E_m(L)$ and the operator ∇ in (80) acts in $E_{\mu}(L)$. Clearly, $r_{m,\mu}^{\triangleright}$ surjectively maps

$$\begin{aligned} \mathfrak{A}_m(L)^{\circ} &\text{ onto } \mathfrak{A}_{\mu}(L)^{\circ}, \mathfrak{A}_m(L)^{\vee} \text{ onto } \mathfrak{A}_{\mu}(L)^{\vee}, \\ \mathfrak{A}_m(L)^{\wedge} &\text{ onto } \mathfrak{A}_{\mu}(L)^{\wedge}, \mathfrak{A}_m(L)^{\circ\wedge} \text{ onto } \mathfrak{A}_{\mu}(L)^{\circ\wedge}, \end{aligned}$$

$\mathfrak{A}_m(L)^{\vee\wedge}$ onto $\mathfrak{A}_\mu(L)^{\vee\wedge}$, $\mathfrak{A}_m(L)^{\circ\rightarrow}$ onto $\mathfrak{A}_\mu(L)^{\circ\rightarrow}$,
 $\mathfrak{A}_m(L)^{\vee\rightarrow}$ onto $\mathfrak{A}_\mu(L)^{\vee\rightarrow}$, $\mathfrak{A}_m(L)^{\circ\wedge\rightarrow}$ onto $\mathfrak{A}_\mu(L)^{\circ\wedge\rightarrow}$,
 $\mathfrak{A}_m(L)^{\vee\wedge\rightarrow}$ onto $\mathfrak{A}_\mu(L)^{\vee\wedge\rightarrow}$. Since the diagram

$$\begin{array}{ccc} E_m(L) & \xrightarrow{r_{m,\mu}} & E_\mu(L) \\ \alpha \downarrow & & \downarrow r_{m,\mu}^\triangleright(\alpha) \\ E_m(L) & \xrightarrow{r_{m,\mu}} & E_\mu(L) \end{array}$$

is commuative and therefore

$$(81) \quad r_{m,\mu}\alpha = r_{m,\mu}^\triangleright(\alpha)r_{m,\mu},$$

it follows that $r_{m,\mu}$ surjectively maps $\text{Ker}(r_{m,\mu}\alpha)$ onto

$$\text{Ker}(r_{m,\mu}^\triangleright(\alpha)) \supset r_{m,\mu}\text{Ker}(\alpha).$$

Lemma 6. *If $\mu \in m - 1 + \mathbb{N}$ and $\alpha \in \mathfrak{A}_m(L)^\wedge$, then the operator α bijectively maps $\text{Ker}(r_{m,\mu})$ onto $\text{Ker}(r_{m,\mu})$.*

Proof. Proof is given in [72], Lemma 3.

Corollary 1. *Let $\mu \in m - 1 + \mathbb{N}$ and let $\alpha \in \mathfrak{A}_m(L)^\wedge$. If*

$$g \in E_m(L), x \in E_\mu(L), m \leq \mu, \alpha \in \mathfrak{A}_m(L)^\wedge,$$

$$r_{m,\mu}(g) = (r_{m,\mu}^\triangleright(\alpha))(x),$$

then there exists a unique $y \in E_m(L)$ such that

$$\alpha(y) = g, r_{m,\mu}(y) = x;$$

Proof. Proof is given in [72], Corollary 1 to the Lemma 3.

Corollary 2. *Let $\mu \in m - 1 + \mathbb{N}$ and $\alpha \in \mathfrak{A}_m(L)^\wedge$. Then $r_{m,\mu}$ bijectively maps $\text{Ker}(\alpha)$ onto $\text{Ker}(r_{m,\mu}^\triangleright(\alpha)) = r_{m,\mu}(\text{Ker}(\alpha))$.*

Proof. Proof is given in [72], Corollary 2 to the Lemma 3.

If for the equation (2) are fulfilled the conditions (1) then

$$(82) \quad a_k = (a_k(0), a_k(1), \dots, a_k(\nu), \dots) \in E_0^\rightarrow(\mathbb{C}),$$

where $k = 0, \dots, n$. Moreover, $a_n = 1_{E_0(\mathbb{C})}$, for α in (79) $\text{Ker}(\alpha)$ coincides with the linear over $\mathbb{C}$ space of all the solutions of the equation (2), polynomial (4) is equal to the polynomial $P(\alpha, z) = P(r_{0,m}^\triangleright(\alpha), z)$, where $m \in \mathbb{N}$, and the set $\text{Ker}(r_{0,m}^\triangleright(\alpha))$ coincides with linear over $\mathbb{C}$ space V_m of all the solutions of the equation (3).

Let $\mathfrak{v}$ be the element in $E_{0,0}$, for which

$$\mathfrak{v}(\nu) = \frac{1}{\nu + 1},$$

where $\nu \in \mathbb{N} - 1$. Clearly, $r_{0,m}(\mathfrak{v})E_{m,\infty}(\mathbb{C}) \subset E_{m,0}(\mathbb{C})$ for any $m \in \mathbb{N} - 1$. Let

$$E_{m,0}^\triangleright(L) = r_{0,m}(\mathfrak{v})E_{m,\infty}(L), E_m^\triangleright(L) = L + E_{m,0}^\triangleright(L).$$

Let us consider the ring $\mathfrak{A}_m^\succ(L)$ generated by the operator ∇ and by all the operators μ_a , where $a \in E_m(L)^\succ$. The Lemma 5 may be reformulated as follows:

Lemma 7. *Let $\alpha \in \mathfrak{A}_0^\succ(\mathbb{C}) \cap \mathfrak{A}_0^\vee(\mathbb{C})$, and $P(\alpha, z)$ coincides with the polynomial $T(z)$ in (4) and (64).*

Then there exist $m \in \mathbb{N}$ and representation of the operator $r_{0,m}^\triangleright(\alpha)$ in the form

$$(83) \quad r_{0,m}^\triangleright(\alpha) = \eta\beta$$

such that

$$(84) \quad \eta \in \mathfrak{A}_m^\succ(\mathbb{C}) \cap \mathfrak{A}_m^\vee(\mathbb{C}), \deg(\eta) = q$$

$$(85) \quad \beta \in \mathfrak{A}_m^\succ(\mathbb{C}) \cap \mathfrak{A}_m^{\vee\wedge}(\mathbb{C}), \deg(\beta) = p = n - q,$$

and the polynomials $P(\beta, z)$, $P(\eta, z)$ coincide respectively with the polynomials $T_1(z)$, $T_2(z)$ in (65).

Lemma 8. *Let are fulfilled all the conditions of the Theorem 7. Let $\alpha \in \mathfrak{A}_0^\succ(\mathbb{C}) \cap \mathfrak{A}_0^\vee(\mathbb{C})$ corresponds to the equation (2), i.e. (with $m = 0$) (79) holds*

$$(86) \quad \alpha = \sum_{k=0}^n \mu_{a_k} \circ \nabla^k,$$

where $a_n = 1_{E_0(\mathbb{C})}$ and

$$(87) \quad a_k = (a_k(0), a_k(1), \dots, a_k(\nu), \dots) \in E_0^\succ(\mathbb{C}),$$

for $k = 0, \dots, n$.

Let the characteristic polynomial (4) is represented as product

$$(88) \quad P(\alpha, z) = T(z) = \prod_{k=1}^{s+1} T_i(z),$$

where

$$(89) \quad T_i(z) = \sum_{\alpha=0}^{e_i} b_{i,\alpha}^\sim z^\alpha,$$

with $b_{i,e_i} = a_n^\sim = 1$ and absolute value of each root of the polynomial $T_i(z)$ is equal to ρ_i .

Then there exist $m \in \mathbb{N}$ and representation of the operator $r_{0,m}^\triangleright(\alpha)$ in the form

$$(90) \quad r_{0,m}^\triangleright(\alpha) = \prod_{i=0}^s \beta_{s+1-i}$$

such that

$$(91) \quad \beta_i \in \mathfrak{A}_m^\succ(\mathbb{C}) \cap \mathfrak{A}_m^\vee(\mathbb{C}), \deg(\beta_i) = e_i,$$

$$P(\beta_i, z) = T_i(z),$$

where $i = 1, \dots, s+1$ and

$$(92) \quad \beta_i \in \mathfrak{A}_m^\wedge(\mathbb{C}),$$

where $i = 1, \dots, s$.

Proof. The assertion of the Lemma may be obtained by means of the sequentially applying of the Lemma 7. ■

Let

$$(93) \quad \beta_\theta^* = \prod_{i=0}^{\theta-1} \beta_{\theta-i}, \quad \beta_\theta^\wedge = \prod_{i=s+1-\theta}^s \beta_{s+1-i},$$

where $\theta = 1, \dots, s+1$. In view of (refeq:2cdj),

$$(94) \quad r_{0,m}^\triangleright(\alpha) = \beta_{\theta+1}^\wedge \beta_\theta^*,$$

$\theta = 1, \dots, s$.

Let $C > 0$, $n \in \mathbb{N}$,

$$(95) \quad w_{C,n}(\nu) = \left(\frac{C}{\nu} \right)^{\nu/n},$$

where $\nu \in \mathbb{N}$; let further $\rho > 0$ and

$$(96) \quad v_{C,n,\rho}(\nu) = \rho^\nu \exp \left(C \left((\nu)^{1-1/n} + \ln(\nu) \right) \right),$$

where $\nu \in \mathbb{N}$.

Let $\theta_1 \in \{1, \dots, s\}$. Replacing m by some bigger m and applying to $\beta_{\theta_1}^\wedge$ in (93) the Theorem 7, we see that there exist $A^\vee > 0$, $m \in \mathbb{N}$, $\alpha^\vee(\nu) > 0$ with $\nu \in m + \mathbb{N} - 1$ and the subspaces $R_{m,\theta_1,\theta_1}^\vee, \dots, R_{m,\theta_1,s+1}^\vee$ of the space $R_m = \text{Ker}(\beta_{\theta_1}^\wedge)$ such that

$$\lim_{\nu \rightarrow \infty} \alpha^\vee(\nu) = 0,$$

$$R_m = R_{m,\theta_1,\theta_1}^\vee \oplus \dots \oplus R_{m,\theta_1,s+1}^\vee, \quad \dim_{\mathbb{C}}(R_{m,\theta_1,i}^\vee) = e_i, \quad \theta_1 \leq i \leq s+1,$$

if

$$(97) \quad r \in R_{m,\theta_1,\theta}^\vee$$

for some $\theta \in \{\theta_1, \dots, s\}$, then

$$(98) \quad \exp(-A^\vee(\ln(\nu) + \nu^{1-1/k_\theta})) \omega_{q,r}(m)(\rho_\theta)^\nu \leq \omega_{q,r}(\nu),$$

where $\nu \in \mathbb{N} + m - 1$; moreover, the spaces

$$R_{m,\theta_1,j}^\wedge = R_{m,\theta_1,j}^\vee \oplus \dots \oplus R_{m,\theta_1,s+1}^\vee,$$

where $j = \theta_1, \dots, s+1$ and natural projections $\pi_{\theta_1,j}^\vee$ of the space $R_{m,\theta_1,j}^\wedge$ onto the space $R_{m,\theta_1,j}^\vee$, where $j = \theta_1, \dots, s$, have the following properties:

if

$$(99) \quad r \in R_{m,\theta_1,\theta}^\wedge$$

for some $\theta \in \{\theta_1, \dots, s\}$, then

$$(100) \quad \omega_{q,r}(\nu) \leq (\rho_\theta)^\nu \exp(A^\vee(\ln(\nu) + \nu^{1-1/k_\theta})) \omega_{q,r}(m),$$

$$(101) \quad \omega_{q,r}(\nu) \geq (\omega_{q,\pi_{\theta_1,\theta}(r)}^\vee(m) - \alpha^\vee(\nu) \omega_{q,r}(m)) \times \\ (\rho_\theta)^\nu \exp(-A^\vee(\ln(\nu) + \nu^{1-1/k_\theta})),$$

where $\nu \in \mathbb{N} + m - 1$; if $e_{s+1} > 0$, and

$$(102) \quad r \in R_{m,\theta_1,s+1}^\vee (= R_{m,\theta_1,s+1}^\wedge),$$

then

$$(103) \quad |r(\nu)| \leq (A^\vee/\nu)^{\nu/k^*} \omega_{q,r}(m),$$

where $\nu \in \mathbb{N} + m - 1$.

Let

$$(104) \quad \beta = \beta_1^* = \beta_1, \eta = \beta_2^\wedge.$$

Then, in view of (94), (83) holds.

Lemma 9. *Let all the conditions of the Theorem 7 are fulfilled. For sufficient big $m \in \mathbb{N}$ there exists the splitting monomorphism*

$$\psi_2^\wedge: Ker(\eta) \rightarrow Ker(\eta\beta)$$

with following properties:

(a)

$$Ker(\beta) = V_{m,1}^\vee, \psi_2^\wedge(Ker(\eta)) = V_{m,2}^\wedge,$$

where $V_{m,1}^\vee$ and $V_{m,2}^\wedge$ are defined in the assertion of the Theorem 7;

(b) the monomorphism $\psi_2^\wedge$ maps isomorphically the space $R_{m,2,j}^\vee$, where $j = 2, \dots, s+1$, onto the space $V_{m,j}^\vee$;

(c) the map $\beta_1 \psi^{wedge}_2$ coincides with identity map $Ker(\eta) \rightarrow Ker(\eta)$;

(d) the restriction of the map $\psi^{wedge}_2 \beta_1$ on the space $V_{m,2}^\wedge$ coincides with the identity map $V_{m,2}^\wedge \rightarrow V_{m,2}^\wedge$;

(e) let I be the identity map $V_m \rightarrow V_m$; then the natural projection π_1 of the space $V_{m,1}^\wedge = V_m$ onto $V_{m,1}^\vee$ coincides with restriction of the map $I - \psi_2^\wedge \beta_1$ on the space $V_{m,1}^\vee$;

(f) if $j \in [2, s] \cap \mathbb{N}$, then natural projection π_j of the space $V_{m,j}^\wedge$ onto the space $V_{m,j}^\vee$ coincides with the restriction of the map $\psi_2^\wedge \pi_{2,j}^\vee \beta_1$ on the space $V_{m,j}^\wedge$ and the projection $\pi_{2,j}^\vee$ coincides with the restriction of the map $\beta_1 \pi_j, \psi_2^\wedge$ on the space $R_{2,j}^\wedge$.

Proof. See the proof of the Theorem 7 in [77], especially the section 2 and section 3. ■.

Remark 2. I use this opportunity to make a corrections in [77]. On the page 13, third line from the bottom must stand $\psi_m \pi_j^\vee \phi_m$ instead $\psi_m \pi_j^\vee$; on the second line from the bottom must stand $\pi_j^\vee = \phi_m \pi_j \psi_m$ instead $\pi_j^\vee = \phi_m \pi_j$.

Lemma 10. *Let all the conditions of the Lemma 9 are fulfilled. Then (for sufficient big m)*

$$(105) \quad V_{m,\theta}^* = Ker(\beta_{1,\theta}^*),$$

where $\theta = 1, \dots, s$.

Proof. We use induction on s . For $s = 1$ or $\theta = 1$ the assertion of the Lemma directly follows from the Lemma 9.

Let $s > 1$ and assertion of the lemma is true for $s - 1$. Let $\theta > 1$. Since the monomorphism $\psi_2^\wedge$ maps isomorphically the space $R_{m,2,j}^\vee$, where $j = 2, \dots, \theta$, onto the space $V_{m,j}^\vee$ and the map $\beta_1 \psi_2^\wedge$ coincides with identity map $\text{Ker}(\eta) \rightarrow \text{Ker}(\eta)$, it follows that the restriction of the map β_1 on the space $V_{m,2}^\vee \oplus \dots \oplus V_{m,\theta}^\vee$ is an isomorphism of $V_{m,2}^\vee \oplus \dots \oplus V_{m,\theta}^\vee$ onto $R_{m,2,\theta}^*$. According to the inductive hypothesis,

$$R_{m,2,\theta}^* = \text{Ker}(\beta_{2,\theta}^*).$$

Therefore, if $y \in V_{m,2}^\vee \oplus \dots \oplus V_{m,\theta}^\vee$ then $\beta_{1,\theta}^*(y) = \beta_{2,\theta}^*(\beta_1(y)) = 0$; moreover, since also $V_{m,1}^\vee = \text{Ker}(\beta_1)$, it follows that $V_{m,\theta}^* \subset \text{Ker}(\beta_{1,\theta}^*)$. Since, according to the Theorem 7,

$$(106) \quad \dim_{\mathbb{C}}(V_{m,\theta}) = \sum_{i=1}^{\theta} e_i =$$

$$\dim_{\mathbb{C}}(\text{Ker}(\beta_{1,\theta}^*)) = n_\theta := \deg(\beta_{1,\theta}^*),$$

it follows that (105) holds. ■

Proof of the Theorem 8. In view of (105), (106), we apply to $\beta_{1,\theta}^*$ the Lemma 3. Since θ plays the role of s in the lemma 3 now and

$$\max(k_1, \dots, k_\theta) \leq k = \max(k_1, \dots, k_s),$$

it follows that there exist $A = A_\theta > 0$, $m \in \mathbb{N}$ such that

$$(107) \quad \exp(-A(\ln(\nu) + \nu^{1-1/k}))(\rho_s)^\nu \omega_{n_\theta,y}(m) \leq$$

$$\omega_{n_\theta,y}(\nu) \leq \omega_{n,y}(\nu)$$

for any $y \in V_{m,\theta}^*$. Since both the functions $y \rightarrow \omega_{n_\theta,y}(m)$, $y \in V_{m,\theta}^*$, and $y \rightarrow \omega_{n,y}(m)$, $y \in V_{m,\theta}^*$ are two norms on the n_θ -dimensional space $V_{m,\theta}^*$ there exists $B = B_\theta > 0$ such that

$$\omega_{n,y}(m) \exp(-B) \leq \omega_{n_\theta,y}(m)$$

for any $y \in V_{m,\theta}^*$. Then

$$(108) \quad \exp(-A(\ln(\nu) + \nu^{1-1/k}) - B)(\rho_\theta)^\nu \omega_{n,y}(m) \leq \omega_{n,y}(\nu)$$

for any $y \in V_{m,\theta}^*$. ■

§3. Proof of the theorem 9.

Theorem 9. Let for some $\theta \in \{1, \dots, s\}$ is given a linear map ξ_θ of the space $V_{m,\theta}^*$ into $V_{m,\theta+1}^\wedge$. Let I_θ^* is the identity map $V_{m,\theta}^* \rightarrow V_{m,\theta}^*$. Then for

$$V = (I_\theta^* + \xi_\theta)(V_{m,\theta}^*)$$

holds the assertion of the Theorem 10.

Proof of the theorem 9. Since $V_{m,\theta}^*$ is finite-dimensional linear space over $\mathbb{C}$, it follows that the map ξ_θ is continuous. Therevore there exists $C > 0$ such that

$$(109) \quad \omega_{n_\theta, \xi_\theta(y)}(m) \leq \exp(C) \omega_{n_\theta, \xi_\theta(y)}(m)$$

for any $y \in V_{m,\theta}^*$. According to the Theorem 7 and Theorem 8, there exist a numbers $A > 0$, $m \in \mathbb{N}$, such that, if $y \in V_{m,\theta}^*$, $\nu \in m - 1 + \mathbb{N}$, then

$$(110) \quad \omega_{n, \xi_\theta(y)}(\nu) \leq \exp(A(\ln(\nu) + \nu^{1-1/k_{\theta+1}}))(\rho_\theta)^\nu \omega_{n, \xi_\theta(y)}(m)$$

for $\theta < s$,

$$(111) \quad \omega_{n_\theta, \xi_\theta(y)}(m) = 0$$

for $\theta = s$ and $k^* = k_{s+1} = 0$,

$$(112) \quad |\xi_\theta(y)(\nu)| \leq (A/\nu)^{\nu/k^*} \omega_{n, \xi_\theta(y)}(m)$$

for $\theta = s$ and $k^* = k_{s+1} > 0$,

$$(113) \quad \exp(-A(\ln(\nu) + \nu^{1-1/k}))(\rho_\theta)^\nu \omega_{n,y}(m) \leq \omega_{n,y}(\nu).$$

Therefore

$$(114) \quad \omega_{n, y+\xi_\theta(y)}(\nu) \geq \omega_{n,y}(\nu) - \omega_{n, \xi_\theta(y)}(\nu) \geq \exp(-A(\ln(\nu) + \nu^{1-1/k}))(\rho_\theta)^\nu \omega_{n,y}(m)$$

for $\theta = s$ and $k^* = k_{s+1} = 0$,

$$(115) \quad \begin{aligned} \omega_{n, y+\xi_\theta(y)}(\nu) &\geq \omega_{n,y}(\nu) - \omega_{n, \xi_\theta(y)}(\nu) \geq \\ &\exp(-A(\ln(\nu) + \nu^{1-1/k}))(\rho_\theta)^\nu \omega_{n,y}(m) - \\ &|\xi_\theta(y)(\nu)| \leq (A/\nu)^{\nu/k^*} \exp(C) \omega_{n,y}(m) = \\ &\exp(-A(\ln(\nu) + \nu^{1-1/k}))(\rho_\theta)^\nu \omega_{n,y}(m) \times \\ &(1 + \exp(A(\ln(\nu) + \nu^{1-1/k}))(\rho_\theta)^{-\nu} (A/\nu)^{\nu/k^*} \exp(C)) \end{aligned}$$

for $\theta = s$ and $k^* = k_{s+1} > 0$,

$$(116) \quad \begin{aligned} \omega_{n, y+\xi_\theta(y)}(\nu) &\geq \omega_{n,y}(\nu) - \omega_{n, \xi_\theta(y)}(\nu) \geq \\ &\exp(-A(\ln(\nu) + \nu^{1-1/k}))(\rho_\theta)^\nu \omega_{n,y}(m) - \\ &(A/\nu)^{\nu/k^*} \exp(C) \omega_{n,y}(m) = \\ &\exp(-A(\ln(\nu) + \nu^{1-1/k}))(\rho_\theta)^\nu \omega_{n,y}(m) \times \\ &(1 - \exp(A(\ln(\nu) + \nu^{1-1/k}))(\rho_\theta)^{-\nu} (A/\nu)^{\nu/k^*} \exp(C)) \end{aligned}$$

for $\theta = s$ and $k^* = k_{s+1} > 0$,

$$(117) \quad \begin{aligned} \omega_{n, y+\xi_\theta(y)}(\nu) &\geq \omega_{n,y}(\nu) - \omega_{n, \xi_\theta(y)}(\nu) \geq \\ &\exp(-A(\ln(\nu) + \nu^{1-1/k}))(\rho_\theta)^\nu \omega_{n,y}(m) - \end{aligned}$$

$$\begin{aligned} & \exp(A(\ln(\nu) + \nu^{1-1/k}))(\rho_\theta)^\nu \exp(C)\omega_{n,y}(m) = \\ & \exp(-A(\ln(\nu) + \nu^{1-1/k}))(\rho_\theta)^\nu \omega_{n,y}(m) \times \\ & (1 - \exp(2A(\ln(\nu) + \nu^{1-1/k}))(\rho_{\theta+1}/\rho_\theta)^\nu \exp(C)) \end{aligned}$$

for $\theta < s$. ■

Remark 3. The value of $\omega_{n,y}(m)$ in (8) may be much bigger than the value of $\omega_{n\pi_\theta(y)}(m)$. Therefore we cannot (as show a simple examples) for fixed value of A in the inequality (11) to replace the linear space V of the Theorem 10 by the set $V_m \setminus V_{m,\theta+1}^\wedge$. But in (116) – (117) $\omega_{n,y}(m)$ is carried out the brackets, and the value in the brackets tends to 1, when $\nu \rightarrow +\infty$, being greater than $1/2$ for $\nu \in m - 1 + \mathbb{N}$, with sufficient big $m \in \mathbb{N}$, which depends only from the equation.

§4. Proof of the theorem 10.

Proof of the theorem 10. Let π^* and $\pi^\wedge$ are restrictions on V of the natural endomorphisms of the space V_m onto respectively $V_{m,\theta}^*$ and $V_{m,\theta+1}^\wedge$ and let I_0 be the identity map $V \rightarrow V$. Then

$$(118) \quad I_0 = \pi^* + \pi^\wedge$$

Since $\text{Ker}(\pi^*) \subset V \cap V_{m,\theta+1}^\wedge = \{0\}$, it follows that π^* is an isomorphism of the space V onto linear subspace V' of the space $V_{m,\theta}^*$. Let τ be the inverse isomorphism to the isomorphism π^* . Then $\pi^*\tau$ is identity map $V' \rightarrow V'$ and it follows from (118) that

$$(119) \quad \tau = \pi^*\tau + \pi^\wedge\tau.$$

Clearly, the linear map $\pi^\wedge\tau : V' \rightarrow V_{m,\theta+1}^\wedge$ have an extension

$$(120) \quad \xi_\theta : V_{m,\theta}^* \rightarrow V_{m,\theta+1}^\wedge.$$

It follows from (119) that

$$V = \tau(V') \subset (I + \xi_\theta)V_{m,\theta}^*,$$

where I is identity map $V_{m,\theta} \rightarrow V_{m,\theta}$ and ξ_θ is a linear map in (120).

So, the Theorem 10 is Corollary of the Theorem 9. ■

§5. The case of the general difference equation of the Poincaré type.

If (12) and (13) hold instead of (1), then making use of the above arguments we obtain the following chainges in the Theorem 7 and theorem 10. For any $\varepsilon > 0$ there exists a constant $A^\wedge = A^\wedge(\varepsilon) > 0$ such that, if $y \in V_{m,\theta}^\wedge$ with some $\theta \in \{1, \dots, s\}$, then (instead of (7) the following inequality holds)

$$(121) \quad \omega_{n,y}(\nu) \leq \exp(A^\wedge(\rho_\theta \exp(\varepsilon)))^\nu \omega_{n,y}(m),$$

where $\nu \in m + \mathbb{N} - 1$; if $y \in V_{m,s+1}^\vee (= V_{m,s+1}^\wedge)$, then (instead of (10) the following inequality holds)

$$(122) \quad \omega_{n,y}(\nu) \leq \exp(A^\wedge(\exp(-\varepsilon)))^\nu \omega_{n,y}(m),$$

where $\nu \in m + \mathbb{N} - 1$.

Let further V be an arbitrary linear subspace of V_m such that

$$V \cap V_{m,\theta+1} = \{0\},$$

where $\theta \in \{1 \dots, s\}$. Then for this subspace V and any $\varepsilon > 0$ there exists a constant $A^\vee = A^\vee(V, \varepsilon) > 0$ such that (instead of (11) the following inequality holds)

$$(123) \quad \exp(-A^\vee)(\rho_\theta \exp(-\varepsilon))^\nu \omega_n(y)(m) \leq \omega_{n,y}(\nu)$$

where $y \in V$ and $\nu \in m + \mathbb{N} - 1$.

Corollary. (See [51], Theorem 3 and [75], Lemma 16). *Let as before (12) and (13) hold instead of (1). Let V be a r -dimensional subspace of V_m , let*

$$V \cap V_{m,s+1}^\vee = \{0\}$$

and let $\{y_1(\nu), \dots, y_r(\nu)\}$ be a basis of the space V . Let

$$k_3(V) = \max\{k \in \mathbb{Z}: 1 \leq k \leq s, V \subset V_{m,k}^\wedge\},$$

and

$$k_4(V) = \min\{k \in \mathbb{Z}: 1 \leq k \leq s, V \cap V_{m,k+1}^\wedge = \{0\}\}.$$

For $X = (x_1, \dots, x_r)$, $X \in \mathbb{C}^r$, let

$$h(X) = \max\{|x_1|, \dots, |x_r|\},$$

$$y = y^\vee(X, \nu) = x_1 y_1^\vee(\nu) + \dots + x_r y_r^\vee(\nu).$$

Then for every $\varepsilon \in (0, 1)$ there exist $C_3(\varepsilon) > 0$ and $C_4(\varepsilon) > 0$ such that

$$C_4(\varepsilon)(\rho_{k_4}(1 - \varepsilon))^\nu h(X) \leq \omega_{n,y}(\nu) \leq C_3(\varepsilon)(\rho_{k_3} + \varepsilon)^\nu h(X).$$

Proof The functions $h(X)$ and the restriction of $\omega_{n,y}(m)$ on V are two norms on the r -dimensional over $\mathbb{C}$ linear space V . Therefore there exists a constant $C_5 > 0$ such that $h(X) \leq C_5 \omega_{n,y}(m)$ and $\omega_{n,y}(m) \leq C_5 h(X)$. The assertion of the Corollary directly follows from (121) and (123) now. ■

References.

- [1] R.Apéry, Interpolation des fractions continues
et irrationalité de certaines constantes,
Bulletin de la section des sciences du C.T.H., 1981, No 3, 37 – 53;
- [2] F.Beukers, A note on the irrationality of $\zeta(2)$ and $\zeta(3)$,
Bull. London Math. Soc., 1979, 11, 268 – 272;
- [3] A.van der Porten, A proof that Euler missed...Apéry's proof of the irrationality of $\zeta(3)$,
Math Intellegencer, 1979, 1, 195 – 203;
- [4] W. Maier, Potenzreihen irrationalen Grenzwertes,
J.reine angew. Math., 156, 1927, 93 – 148;
- [5] E.M. Nikišin, On irrationality of the values of the functions $F(x,s)$ (in Russian),
Mat.Sb. 109 (1979), 410 – 417;
English transl. in Math. USSR Sb. 37 (1980), 381 – 388;
- [6] G.V. Chudnovsky, Pade approximations to the generalized hyper-geometric functions
I,J.Math.Pures Appl., 58, 1979, 445 – 476;
- [7] —————, Transcendental numbers, Number Theory,Carbondale,

- [8] —————, Lecture Notes in Math, Springer-Verlag, 1979, 751, 45 – 69;
- [9] —————, Approximations rationnelles des logarithmes de nombres rationelles
C.R.Acad.Sc. Paris, Série A, 1979, 288, 607 – 609;
- [10] —————, Formules d'Hermite pour les approximants de Padé de logarithmes
et de fonctions binômes, et mesures d'irrationalité,
C.R.Acad.Sc. Paris, Série A, 1979, t.288, 965 – 967;
- [11] —————, Un système explicite d'approximants de Padé
pour les fonctions hypergéométriques généralisées,
avec applications à l'arithmétique,
C.R.Acad.Sc. Paris, Série A, 1979, t.288, 1001 – 1004;
- [12] —————, Recurrences defining Rational Approximations
to the irrational numbers, Proceedings
of the Japan Academie, Ser. A, 1982, 58, 129 – 133;
- [13] —————, On the method of Thue-Siegel,
Annals of Mathematics, 117 (1983), 325 – 382;
- [14] K.Alladi and M. Robinson, Legendre polynomials and irrationality,
J. Reine Angew.Math., 1980, 318, 137 – 155;
- [15] A. Dubitskas, An approximation of logarithms of some numbers,
Diophantine approximations II, Moscow, 1986, 20 – 34;
- [16] —————, On approximation of $\pi/\sqrt{3}$ by rational fractions,
Vestnik MGU, series 1, 1987, 6, 73 – 76;
- [17] S.Eckmann, Über die lineare Unabhängigkeit der Werte gewisser Reihen,
Results in Mathematics, 11, 1987, 7 – 43;
- [18] M.Hata, Legendre type polynomials and irrationality measures,
J. Reine Angew. Math., 1990, 407, 99 – 125;
- [19] A.O. Gelfond, Transcendental and algebraic numbers (in Russian),
GIT-TL, Moscow, 1952;
- [20] H.Bateman and A.Erdélyi, Higher transcendental functions, 1953,
New-York – Toronto – London, Mc. Grow-Hill Book Company, Inc.;
- [21] E.C.Titchmarsh, The Theory of Functions, 1939, Oxford University Press;
- [22] E.T.Whittaker and G.N. Watson, A course of modern analysis,
1927, Cambridge University Press;
- [23] O.Perron, Über die Poincaresche Differenzengleichung,
Journal für die reine und angewandte mathematik,
1910, 137, 6 – 64;
- [24] A.O.Gelfond, Differenzenrechnung (in Russian), 1967, Nauka, Moscow.
- [25] A.O.Gelfond and I.M.Kubenskaya, On the theorem of Perron
in the theory of difference equations (in Russian),
IAN USSR, math. ser., 1953, 17, 2, 83 – 86.
- [26] M.A.Evgrafov, New proof of the theorem of Perron
(in Russian), IAN USSR, math. ser., 1953, 17, 2, 77 – 82;
- [27] G.A.Frejman, On theorems of Poincaré and Perron
(in Russian), UMN, 1957, 12, 3 (75), 243 – 245;
- [28] N.E.Nörlund, Differenzenrechnung, Berlin, Springer Verlag, 1924;
- [29] I.M.Vinogradov, Foundations of the Number Theory, (in Russian), 1952, GIT-TL;
- [30] —————, J. Diedonne, Foundations of modern analysis,
Institut des Hautes Études Scientifiques, Paris,
Academic Press, New York and London, 1960
- [31] CH.-J. de la Vallée Poussin, Course d'analyse infinitésimale,
Russian translation by G.M.Fikhtengolts,
GT-TI, 1933;
- [32] H.WEYL, Algebraic theory of numbers,
1940,
Russian translation by L.I.Kopejkina;
- [33] G.Rhin and C. Viola.
On a permutation group related to $\zeta(2)$,
Acta Arithmetica 77(1996), 25 – 56;
- [34] G.Rhin and C. Viola. The group structure for $\zeta(3)$,
Acta Arithmetica 97(2001), 269 – 293;

- [34] M. Hata. A new irrationality measure for $\zeta(3)$,
Acta Arithmetica 92(2000), 47 – 57;
- [35] T.Rivual. La fonction zêta de Riemann prend une infinité
valeurs irrationnelles aux entiers impairs,
C.R.Acad.Sci.Paris, série 1, p. 267 – 270, 2000.
- [36] K.Boll, T.Rivual. Irrationalité d’une infinité
valeurs la fonction zêta aux entiers impairs,
Invent. math., 146, 193 – 207 (2001);
- [37] W.Zudilin. One from the numbers $\zeta(5)$, $\zeta(7)$, $\zeta(9)$, $\zeta(11)$ is irrational.
(in Russian) UMN 56(4) (2001), 149 – 150;
- [38] W.Zudilin. Apéry’s Theorem and problems for the values
of ζ -function of Riemann and their q -analogs.
(in Russian) Moscow state university, doctoral thesis,
Moscow 2004;
- [38] L.A.Gutnik, On the decomposition of the difference operators of Poincaré type
(in Russian), VINITI, Moscow, 1992, 2468 – 92, 1 – 55;
- [39] ———, On the decomposition of the difference operators
of Poincaré type in Banach algebras
(in Russian), VINITI, Moscow, 1992, 3443 – 92, 1 – 36;
- [40] ———, On the difference equations of Poincaré type
(in Russian), VINITI, Moscow 1993, 443 – B93, 1 – 41;
- [41] ———, On the difference equations of Poincaré type in normed algebras
(in Russian), VINITI, Moscow, 1994, 668 – B94, 1 – 44;
- [42] ———, On the decomposition of the difference equations of Poincaré type
(in Russian), VINITI, Moscow, 1997, 2062 – B97, 1 – 41;
- [43] ———, The difference equations of Poincaré type
with characteristic polynomial having roots equal to zero
(in Russian), VINITI, Moscow, 1997, 2418 – 97, 1 – 20;
- [44] ———, On the behavior of solutions
of difference equations of Poincaré type
(in Russian), VINITI, Moscow, 1997, 3384 – B97, 1 – 41;
- [45] ———, On the variability of solutions of difference equations of Poincaré type
(in Russian), VINITI, Moscow, 1999, 361 – B99, 1 – 9;
- [46] ———, To the question of the variability of solutions
of difference equations of Poincaré type (in Russian),
VINITI, Moscow, 2000, 2416 – B00, 1 – 22;
- [47] ———, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$,
Max-Planck-Institut für Mathematik,
Bonn, Preprint Series, 2000, 3, 1 – 13;
- [48] ———, On the Irrationality of Some Quantities Containing $\zeta(3)$ (in Russian),
Uspekhi Mat. Nauk, 1979, 34, 3(207), 190;
- [49] ———, On the Irrationality of Some Quantities Containing $\zeta(3)$,
Eleven papers translated from the Russian,
American Mathematical Society, 1988, 140, 45 – 56;
- [50] ———, Linear independence over $\mathbb{Q}$ of dilogarithms at rational points
(in Russian), UMN, 37 (1982), 179-180;
english transl. in Russ. Math. surveys 37 (1982), 176-177;
- [51] ———, On a measure of the irrationality of dilogarithms at rational points
(in Russian), VINITI, 1984, 4345-84, 1 – 74;
- [52] ———, To the question of the smallness of some linear forms
(in Russian), VINITI, 1993, 2413-B93, 1 – 94;
- [53] ———, About linear forms, whose coefficients are logarithms
of algebraic numbers (in Russian),
VINITI, 1995, 135-B95, 1 – 149;
- [54] ———, About systems of vectors, whose coordinates
are linear combinations of logarithms of algebraic numbers
with algebraic coefficients (in Russian),
VINITI, 1994, 3122-B94, 1 – 158;
- [55] ———, On the linear forms, whose

- coefficients are $\mathbb{A}$ - linear combinations
of logarithms of $\mathbb{A}$ - numbers,
VINITI, 1996, 1617-B96, pp. 1 – 23.
- [56] ———, On systems of linear forms, whose
coefficients are $\mathbb{A}$ - linear combinations
of logarithms of $\mathbb{A}$ - numbers,
VINITI, 1996, 2663-B96, pp. 1 – 18.
- [57] ———, About linear forms, whose coefficients
are $\mathbb{Q}$ -proportional to the number $\log 2$, and the values
of $\zeta(s)$ for integer s (in Russian),
VINITI, 1996, 3258-B96, 1 – 70;
- [58] ———, The lower estimate for some linear forms,
coefficients of which are proportional to the values
of $\zeta(s)$ for integer s (in Russian),
VINITI, 1997, 3072-B97, 1 – 77;
- [59] ———, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2000, 3, 1 – 13;
- [60] ———, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$
(the detailed version, part 1), Max-Plank-Institut für Mathematik,
Bonn, Preprint Series, 2001, 15, 1 – 20;
- [61] ———, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$
(the detailed version, part 2), Max-Plank-Institut für Mathematik,
Bonn, Preprint Series, 2001, 104, 1 – 36;
- [62] ———, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$
(the detailed version, part 3), Max-Plank-Institut für Mathematik,
Bonn, Preprint Series, 2002, 57, 1 – 33;
- [63] ———, On the rank over $\mathbb{Q}$ of some real matrices (in Russian),
VINITI, 1984, 5736-84; 1 – 29;
- [64] ———, On the rank over $\mathbb{Q}$ of some real matrices,
Max-Plank-Institut für Mathematik,
Bonn, Preprint Series, 2002, 27, 1 – 32;
- [65] ———, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$
(the detailed version, part 4), Max-Plank-Institut für Mathematik,
Bonn, Preprint Series, 2002, 142, 1 – 27;
- [66] ———, On the dimension of some linear spaces
over finite extension of $\mathbb{Q}$ (part 2),
Max-Plank-Institut für Mathematik, Bonn, Preprint Series,
2002, 107, 1 – 37;
- [67] ———, On the dimension of some linear spaces over $\mathbb{Q}$ (part 3),
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2003, 16, 1 – 45.
- [68] ———, On the difference equation of Poincaré type (Part 1).
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2003, 52, 1 – 44.
- [69] ———, On the dimension of some linear spaces over $\mathbb{Q}$, (part 4)
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2003, 73, 1 – 38.
- [70] ———, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$
(the detailed version, part 5),
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2003, 83, 1 – 13.
- [71] ———, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$
(the detailed version, part 6),
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2003, 99, 1 – 33.
- [72] ———, On the difference equation of Poincaré type (Part 2).
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2003, 107, 1 – 25.

- [73] ———, On the asymptotic behavior of solutions
of difference equation (in English).
Chebyshevskij sbornik, 2003, v.4, issue 2, 142 – 153.
- [74] ———, On linear combinations of logarithms
of algebraic numbers with algebraic coefficients.
International conference
"Diophantine analysis, uniform distributions
and applications"
August 25-30, 2003, Minsk, Belarus
Abstracts
pp. 16 – 17.
- [75] ———, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$,
Bonner Mathematische Schriften Nr. 360,
Bonn, 2003, 360.
- [76] ———, On linear forms with coefficients in $\mathbb{N}\zeta(1 + \mathbb{N})$
(the detailed version, part 7),
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2004, 88, 1 – 27.
- [77] ———, On the difference equation of Poincaré type (Part 3).
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2004, 09, 1 – 34.
- [78] ———, On the dimension of some linear spaces over $\mathbb{Q}$, (part 5)
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2004, 1 – 42.

E-mail: gutnik@gutnik.mccme.ru